

AD-A145 458

NAVAL POSTGRADUATE SCHOOL

Monterey, California





CERENKOV RADIATION IN THE NEIGHBORHOOD OF THE EMISSION THRESHOLD

Fred R. Buskirk and John R. Neighbours

Revised August 1984

Technical Report

Approved for public release; distribution unlimited

Prepared for: Naval Surface Weapons Center R401

84 09 04 050

DTIC FILE COPY

NAVAL POSTGRADUATE SCHOOL Monterey, California

Commodore R. H. Shumaker Superintendent

David Schrady Provost

The work reported herein was supported by the Chief of Naval Research.

Reproduction of all or part of this report is authorized.

This report was prepared by:

F. R. Buskirk

Professor of Physics

J. R. Neighbours

Professor of Physics

Reviewed by:

G. E. Schacher, Chairman Department of Physics J. N. Dyer Dean of Science and Engineering

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
NPS-61-84-007	AD-A145458		
4. TITLE (and Substitle)		3. TYPE OF REPORT & PERSON COVERED	
Cerenkov Radiation in the no of the Emission Threshold	eldupot nood	Technical Report	
Of the purposes surcomes-		S. PERFORMING ORG. REPORT HUMBER	
7. AUTHOR(s) Prod P. Buckirk and		B. CONTRACT OF GRANT NUMBER(s)	
Fred R. Buskirk and John R. Neighbours			
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & BORK UNIT HUMBERS	
Naval Postgraduate School Monterey, California 93943		l	
Monterey, California 93943			
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE	
Naval Surface Weapons Cente	r	Revised August 1984	
N401, White Oak, MD 20910	!	13. NUMBER OF PAGES	
14. MONITORING AGENCY NAME & ADDRESS/II dilloren	i trem Controlling Office)	18. SECURITY CLASS. (of the report)	
		unclassified	
		15a. DECLASSIFICATION/DOWNGRADING	
ł	l	SCHEDULE.	
16. DISTRIBUTION STATEMENT (of this Report)	-		
approved for public release	, distribution	n unlimited	
Supersedea AD-A143088			
17. DISTRIBUTION STATEMENT (of the obstract entered in Block 26, if different from Report)			
16. SUPPLEMENTARY NOTES			
10. SUPPLEMENTARY ROTES			
19. KEY WORDS (Continue on reverse side if necessary an	d (dentify by block member)		
Cerenkov Radiation		-	
Relativistic Electron Beams	;		
20. ABSTRACT (Continue on reverse side II necessary and	I Identify by block master)	· · · · · · · · · · · · · · · · · · ·	
Cerenkov radiation for constant velocity electrons in an infi-			
nite uniform dielectric has a sharp threshold for v (electron) larger than the speed of light in the dielectric. A medium			
lof finite length produces diffraction which smears the			
Crenkov emission angle and lowers the threshold velocity for			
emission.		•	
•			

CERENKOV RADIATION IN THE NEIGHBORHOOD

OF THE EMISSION THERESHOLD

F.R. Buskirk and John R. Neighbours
Physics Department
Naval Postgraduate School

Introduction - Microwave Cerenkov

Cerenkov radiation in the simplest form occurs when a charged particle in uniform motion exceeds the velocity of light in an infinite medium. The radiation is emitted in a come, with the rays occurring at a sharo angle given by $\cos\theta_{\rm C}=c$ (medium)/v (particle), and for an infinite medium and uniform velocity, the radiation would disappear for v < c. The result above is the consequence of requiring the phase of the radiation, emitted at an angle $\theta_{\rm C}$, to remain in phase with the charge as it moves in time. We have studied microwave Cerenkov radiation experimentally and theoretically 1,2,3. These studies describe the microwave or other R.F. emission which is significant for bunches of electrons with dimensions shorter than the wavelength of emitted radiation so that all electrons in the bunch radiate coherently. These effects will be explored elsewhere.

Diffraction Effects

The point of this paper is as follows: If the ideal conditions (constant electron velocity or infinite medium) are changed, the radiation changes, possible dramatically. In ¹ it was noted that for a <u>finite</u> length of medium, diffraction occurred and the Cerenkov angle is smeared. In ^{2,3} these effects were considered further and, besides the smearing of the emission angle,

it was noted that the radiated power may be larger for a finite medium.

The theory is based on Ref. 1, Eq. (A13), which gives the energy radiated per unit solid angle in the frequency range $d\omega$:

$$W (\omega, \vec{k}) d\omega = \frac{1}{16\pi^3} \frac{\mu}{c} \omega^2 \sin^2\theta^2 \left(\frac{L}{v}\right)^2 \frac{\sin^2\theta}{u^2} q^2 F(\vec{k})$$
 (1)

where u is defined below, L is the length of the medium, \vec{k} is the wave number of the emitted radiation, and F is the form factor for the bunch.

Threshold of Cerenkov Radiation

The mechanism allowing the smearing of the angle and the increase in power for a finite medium is relaxation of the phase matching between the electron and the wave. If the wave is emitted at an angle $9 \neq 9_{\rm C}$, the electron and wave will be only slightly out of phase at the end of a finite path L. In fact from 2, the null of the radiation pattern occurs for

$$u = \frac{kL}{2} \left(\frac{c}{-\cos \theta} \right) = \pi \tag{2}$$

Thus we have radiation from θ = 0, to $\theta_{\rm C}$ (where u = 0) and beyond, to $\theta_{\rm R}$ (where u = π). Now note that, if v < c, there is no Cerenkov angle (cos $\theta_{\rm C}$ = c/v has no solution) but $\theta_{\rm R}$ may exist, and radiation occurs below the usually accepted threshold.

This effect was investigated in the optical region 4,5,6 , both theoretically and experimentally by Kobzev and Frank. We may calculate how much the usual Cerenkov threshold could be lowered by noting that the radiation will disappear for all practical purposes when v decreases such that $\theta_{\rm n}$ + o. A simple calculation gives:

$$\frac{1}{2\gamma^2} \text{ (threshold)} = \frac{\lambda}{L} + \Delta, \tag{3}$$

where $1/1-\Delta$ is the relative index of refraction.

Numerical Example

Example: $\Delta = 2.68 \times 10^{-4}$ for air. Let kL = ∞ . Then $\gamma = 43$. Now let L = lm, $\lambda = 1$ cm. Then $\gamma = 7.07$. The change in threshold is indeed dramatic. No attempt has been made to include effects other than abrupt termination of the ideal, infinite uniform medium. This could be accomplished by abruptly stopping the beam, which is assumed to have constant velocity before the stop.

REFERENCES

- 1. F. R. Buskirk and J. R. Neighbours, Physics Rev. A 28, 1531 (1983).
- J. R. Neighbours and F. R. Buskirk (To be published in Phys. Rev. A 29, June (1984).
- 3. J. R. Neighbours and F. R. Buskirk, "Diffraction Effects in Cerenkov Radiation" NPS Technical Report, (1983).
- 4. A. P. Kobzev, Yad. Fiz. 27 1256 (1978) [Sov. J. Nucl. Phys. 27 664 (1978)].
- 5. A. P. Robzev and I. M. Frank Yad. Fiz. 31 1253 (1982) [Sov. J. Nucl. Phys. 31 647, (1980)].
- A. P. Kobzev and I. M. Frank, Yad. Fiz. 34 125 [Sov. J. Nucl. Phys. 34 71 (1981)].



(0)

aebo




Figure Caption:

Qualitative illustration of diffraction effects in Cerenkov radiation associated with a finite length of path. In the upper curve $\mathbf{v} > \mathbf{c}$ and the radiation is spread about the Cerenkov angle (shaded area). In the lower curve, $\mathbf{v} < \mathbf{c}$ but the same diffraction function allows radiation of occur.

DISTRIBUTION LIST

Office of Naval Research CDR R. Swafford 800 N. Quincy Street Arlington, VA 22217	1
Office of Naval Research CDR James Offutt 1030 East Green Street Pasadena, CA 91106	. 1
Library Code 0142 Naval Postgraduate School Monterey, CA 93943	2
Office of Researach Administration Code 012A Naval Postgraduate School Monterey, CA 93943	2
F. R. Buskirk & J. R. Neighbours Naval Postgraduate School Physics Department, Code 61 Monterey, CA 93943	20
Dr. Thomas Starke M4, M.S. P942 Los Alamos National Laboratory Los Alamos, NM 87545	1
MAJ E. W. Poque M4, M.S. P942 Los Alamos National Laboratory Los Alamos, NM 87545	1
Dr. Charles Bowman D442 Los Alamos National Laboratory Los Alamos, NM 87545	1
Dr. Thomas Fessenden L-436 Lawrence Livermore National Laboratorv Box 809 Livermore, CA 94550	1

Dr. Richard Briggs	2
L-321	
Lawrence Livermore National Laboratory	
Box 808	
Livermore, CA 94550	
Dr. C. M. Huddleston	3
R-401	
Naval Surface Weapons Center	
White Oak	
Silver Spring, MD 20910	
CAPT R. L. Topping	1
PMS 405	
Naval Sea Systems Command	
Washington, D.C. 20376	
Dr. David Merritt	1
PMS 405	
Naval Sea Systems Command	
Washington, D.C. 20376	
CDR William Bassett	1
PMS 405	
Naval Sea Systems Command	
Washington, D.C. 20376	
Director, Defense Advanced Research	2
Project Agency	
ATTN: LCOL Richard A. Gullickson	
1400 Wilson Blvd.	
Arlington, VA 22209	
Defense Technical Information Center	2
Cameron Station	
Alexandria, Virginia 22314	